

Polarizing mechanisms for stored p and $\bar{p}$ beams interacting with a polarized target

A.I. Milstein* and V.M.Strakhovenko†

Budker Institute of Nuclear Physics, 630090 Novosibirsk, Russia

(Dated: February 2, 2008)

Kinetics of the polarization buildup at the interaction of stored protons (antiprotons) with a polarized target is considered. It is demonstrated that for small scattering angles, when a projectile remains in the beam, the polarization buildup is completely due to the spin-flip transitions. The corresponding cross sections turn out to be negligibly small for a hydrogen gas target as well as for a pure electron target. For the latter, the filtering mechanism also does not provide a noticeable beam polarization.

PACS numbers: 29.20.Dh, 29.25.Pj, 29.27.Hj

I. INTRODUCTION

At present, the use of a polarized hydrogen gas target to polarize stored antiprotons is widely discussed (see, e.g., [1] and literature therein). The modifications of the method first proposed in [2] are considered. The original idea of spin filtering is based on the dependence of the scattering cross section on the mutual orientation of the target and projectile proton spins. Then the protons having opposite spin projections on the direction of the target polarization go out from the beam with the different rates. As a result, the beam acquires a polarization.

The method suggested in [2] has been realized in the experiment [3], where 23-MeV stored protons scattered on an internal gas target of the polarized hydrogen atoms. In 90 minutes, the intensity of the beam was 5% of the initial one and the polarization degree amounted to 2.4%. In [3], the rate of polarization buildup due to filtering mechanism was also estimated theoretically, taking into account only strong pp interaction. This estimation is noticeably different from the experimental result. The explanation of this disagreement is proposed in [4]. In that paper the importance of the interference of the Coulomb amplitude and the spin-dependent part of the hadronic amplitude is emphasized. Under conditions of the experiment [3], this effect diminishes the corresponding cross section more than by 40 %, thereby, improving essentially agreement with the experiment. Our calculations confirm the estimation of this effect obtained in [4].

The remaining difference between the experimental and theoretical results, which is rather small, is explained in [4] by two new mechanisms. Both mechanisms are related to scattering at the angle ϑ smaller than the acceptance angle $\theta_{acc} \ll 1$, where protons remain in the beam. The first effect, suggested in [5], is due to the interaction of a projectile with the polarized electrons of the hydrogen gas target. The second one, considered in [4], is due to scattering on polarized protons of the target at $\vartheta < \theta_{acc}$. The estimation of these two effects, made in [4], gives the contributions to the rate of polarization buildup close in the absolute values but of the opposite sign. The magnitude of each effect is comparable with that corresponding to filtering effect. The result accounting for all three contributions agrees very well with the experiment. Basing on the results of [5], a method to polarize an antiproton beam was suggested in [1]. The idea of Ref. [1] is in the use of a hydrogen gas target with the high electron and low proton polarizations.

In the present paper, we demonstrate that the consideration of both new effects performed in [4, 5] is not correct. We show that, for scattering at $\vartheta < \theta_{acc}$ (a projectile remains in the beam), the polarization buildup is completely due to spin-flip transitions. For $\vartheta \ll 1$, a noticeable contribution to the spin-dependent part of the differential cross section appears as a result of interference between the spin-independent Coulomb amplitude and spin-dependent amplitude. This is because of the singularity of the Coulomb amplitude at small scattering angles ($\propto 1/\vartheta^2$). Evidently, such interference is absent in scattering with the spin flip of a projectile, and the corresponding cross section is negligibly small. The formulas used in [4, 5] for the description of their two effects correspond to such interference and, therefore, are irrelevant to the kinetics of polarization.

*Electronic address: A.I.Milstein@inp.nsk.su

†Electronic address: V.M.Strakhovenko@inp.nsk.su

II. KINETICS OF POLARIZATION

We consider a beam of particles with the densities in the momentum space $f_+(\mathbf{p}, t)$ and $f_-(\mathbf{p}, t)$, where subscripts correspond to the spin projections $\pm 1/2$ on a quantization axis. Let $W_{\sigma_f \sigma_i}(\mathbf{p}_f, \mathbf{p}_i)$ be the probability of the transition from the state with the momentum $\mathbf{p}_i$ and polarization σ_i to the state with the momentum $\mathbf{p}_f$ and polarization σ_f . Then we have for the densities $f_{\pm}$ the conventional kinetic equation:

$$\begin{aligned} \frac{\partial}{\partial t} f_+(\mathbf{p}, t) &= - \int d\mathbf{p}' [W_{++}(\mathbf{p}', \mathbf{p}) + W_{-+}(\mathbf{p}', \mathbf{p})] f_+(\mathbf{p}, t) \\ &\quad + \int_{\mathbf{p}' \in \Gamma} d\mathbf{p}' [W_{++}(\mathbf{p}, \mathbf{p}') f_+(\mathbf{p}', t) + W_{+-}(\mathbf{p}, \mathbf{p}') f_-(\mathbf{p}', t)], \\ \frac{\partial}{\partial t} f_-(\mathbf{p}, t) &= - \int d\mathbf{p}' [W_{--}(\mathbf{p}', \mathbf{p}) + W_{+-}(\mathbf{p}', \mathbf{p})] f_-(\mathbf{p}, t) \\ &\quad + \int_{\mathbf{p}' \in \Gamma} d\mathbf{p}' [W_{-+}(\mathbf{p}, \mathbf{p}') f_+(\mathbf{p}', t) + W_{--}(\mathbf{p}, \mathbf{p}') f_-(\mathbf{p}', t)]. \end{aligned} \quad (1)$$

Here $\mathbf{p}' \in \Gamma$ means that the momentum $\mathbf{p}'$ belongs to the beam momentum space (the angle between the momentum $\mathbf{p}'$ and the beam axis is less than θ_{acc}). Taking in Eq.(1) the integral over $\mathbf{p} \in \Gamma$, we obtain

$$\begin{aligned} \frac{\partial}{\partial t} \int_{\mathbf{p} \in \Gamma} d\mathbf{p} [f_+(\mathbf{p}, t) - f_-(\mathbf{p}, t)] &= - \int_{\mathbf{p}' \notin \Gamma} d\mathbf{p}' \int_{\mathbf{p} \in \Gamma} d\mathbf{p} [W_{++}(\mathbf{p}', \mathbf{p}) + W_{-+}(\mathbf{p}', \mathbf{p})] f_+(\mathbf{p}, t) \\ &\quad + \int_{\mathbf{p}' \notin \Gamma} d\mathbf{p}' \int_{\mathbf{p} \in \Gamma} d\mathbf{p} [W_{--}(\mathbf{p}', \mathbf{p}) + W_{+-}(\mathbf{p}', \mathbf{p})] f_-(\mathbf{p}, t) \\ &\quad + 2 \int_{\mathbf{p}' \in \Gamma} d\mathbf{p}' \int_{\mathbf{p} \in \Gamma} d\mathbf{p} [W_{+-}(\mathbf{p}', \mathbf{p}) f_-(\mathbf{p}, t) - W_{-+}(\mathbf{p}', \mathbf{p}) f_+(\mathbf{p}, t)], \\ \frac{\partial}{\partial t} \int_{\mathbf{p} \in \Gamma} d\mathbf{p} [f_+(\mathbf{p}, t) + f_-(\mathbf{p}, t)] &= - \int_{\mathbf{p}' \notin \Gamma} d\mathbf{p}' \int_{\mathbf{p} \in \Gamma} d\mathbf{p} [W_{++}(\mathbf{p}', \mathbf{p}) + W_{-+}(\mathbf{p}', \mathbf{p})] f_+(\mathbf{p}, t) \\ &\quad - \int_{\mathbf{p}' \notin \Gamma} d\mathbf{p}' \int_{\mathbf{p} \in \Gamma} d\mathbf{p} [W_{--}(\mathbf{p}', \mathbf{p}) + W_{+-}(\mathbf{p}', \mathbf{p})] f_-(\mathbf{p}, t). \end{aligned} \quad (2)$$

As seen from Eq.(2), in the term where both momenta $\mathbf{p}$ and $\mathbf{p}'$ belong to Γ , only the spin-flip probabilities are present. In other words, scattering without loss of particles may lead to a beam polarization solely due to the spin-flip transitions. Due to phase space cooling, the distributions $f_{\sigma}(\mathbf{p}, t)$ are peaked in the narrow region around a momentum $\mathbf{p}_0$. Then we obtain

$$\begin{aligned} \frac{d}{dt} [N_+(t) - N_-(t)] &= -\Omega_+^{out} N_+(t) + \Omega_-^{out} N_-(t) + 2[\Omega_{+-} N_-(t) - \Omega_{-+} N_+(t)], \\ \frac{d}{dt} [N_+(t) + N_-(t)] &= -[\Omega_+^{out} N_+(t) + \Omega_-^{out} N_-(t)]. \end{aligned} \quad (3)$$

Here

$$\begin{aligned} \Omega_+^{out} &= \int_{\mathbf{p}' \notin \Gamma} d\mathbf{p}' [W_{++}(\mathbf{p}', \mathbf{p}_0) + W_{-+}(\mathbf{p}', \mathbf{p}_0)], \quad \Omega_-^{out} = \int_{\mathbf{p}' \notin \Gamma} d\mathbf{p}' [W_{--}(\mathbf{p}', \mathbf{p}_0) + W_{+-}(\mathbf{p}', \mathbf{p}_0)], \\ \Omega_{+-} &= \int_{\mathbf{p}' \in \Gamma} d\mathbf{p}' W_{+-}(\mathbf{p}', \mathbf{p}_0), \quad \Omega_{-+} = \int_{\mathbf{p}' \in \Gamma} d\mathbf{p}' W_{-+}(\mathbf{p}', \mathbf{p}_0), \quad N_{\pm}(t) = \int_{\mathbf{p} \in \Gamma} d\mathbf{p} f_{\pm}(\mathbf{p}, t). \end{aligned} \quad (4)$$

The solution to Eq.(3), with the initial condition $N_{\pm}(0) = N_0/2$, reads

$$N(t) = N_+(t) + N_-(t) = N_0 \left[\cosh(\Omega t) + \frac{\Omega_{+-} + \Omega_{-+}}{2\Omega} \sinh(\Omega t) \right] \exp(-\Omega_{tot} t),$$

$$P_B(t) = \frac{N_+(t) - N_-(t)}{N_+(t) + N_-(t)} = \frac{\Omega_{+-} - \Omega_{-+} + \frac{1}{2}(\Omega_-^{out} - \Omega_+^{out})}{\Omega + \frac{1}{2}(\Omega_{+-} + \Omega_{-+}) \tanh(\Omega t)}, \quad (5)$$

where $N(t)$ is the total number of particles in the beam, $P_B(t)$ is the beam polarization, and

$$\Omega_{tot} = \frac{1}{2} [\Omega_+^{out} + \Omega_-^{out} + \Omega_{+-} + \Omega_{-+}] ,$$

$$\Omega = \frac{1}{2} \left[(\Omega_-^{out} - \Omega_+^{out})^2 + (\Omega_{+-} + \Omega_{-+})^2 + 2(\Omega_-^{out} - \Omega_+^{out})(\Omega_{+-} - \Omega_{-+}) \right]^{1/2}. \quad (6)$$

For the scattering angle $\vartheta > \theta_{acc}$, the momentum transfer is much larger than $1/a_0$ (a_0 is the Bohr radius, $\hbar = c = 1$). In this case scattering off hydrogen atom can be considered as independent scattering off a free electron and a free proton. The maximum scattering angle of a proton on an electron at rest is less than θ_{acc} for any storage ring. Therefore, scattering on electrons at rest does not contribute to $\Omega_{\pm}^{out}$. If the proton polarization of the hydrogen target is absent (as in the scheme considered in Ref.[1]), then $\Omega_-^{out} = \Omega_+^{out}$ and we have from Eq.(5)

$$P_B(t) = \frac{\Omega_{+-} - \Omega_{-+}}{\Omega_{+-} + \Omega_{-+}} \left[1 - \exp[-(\Omega_{+-} + \Omega_{-+}) t] \right]. \quad (7)$$

If the proton polarization of the target is not small, then $|\Omega_-^{out} - \Omega_+^{out}|$ is much larger than Ω_{+-} and Ω_{-+} . In this case

$$P_B(t) = \tanh \left[\frac{t}{2} (\Omega_-^{out} - \Omega_+^{out}) \right]. \quad (8)$$

III. PROBABILITIES AND CROSS SECTIONS FOR PP SCATTERING

Let us direct the polar axis along the unit vector $\boldsymbol{\nu} = \mathbf{p}_0/p_0$. Then the cross section of pp scattering integrated over the azimuth angle and summed up over the final polarization of both protons in the center-of-mass frame reads

$$d\sigma = 2\pi \sin \vartheta d\vartheta \{ F_0(\vartheta) + (\boldsymbol{\zeta}_t \cdot \boldsymbol{\zeta}_b) F_1(\vartheta) + (\boldsymbol{\zeta}_t \cdot \boldsymbol{\nu})(\boldsymbol{\zeta}_b \cdot \boldsymbol{\nu}) [F_2(\vartheta) - F_1(\vartheta)] \}, \quad (9)$$

where $\boldsymbol{\zeta}_t$ and $\boldsymbol{\zeta}_b$ are the unit polarization vectors of the protons from the target and the beam, respectively. The functions $F_0(\vartheta)$, $F_1(\vartheta)$, and $F_2(\vartheta)$ are

$$\begin{aligned} F_0(\vartheta) &= I_{0000} = \frac{1}{2} \{ |M_1|^2 + |M_2|^2 + |M_3|^2 + |M_4|^2 + 4|M_5|^2 \}, \\ F_1(\vartheta) &= \frac{1}{2} I_{0000} \{ A_{00nn} + \cos^2(\vartheta/2) A_{00mm} + \sin^2(\vartheta/2) A_{00ll} + \sin(\vartheta) A_{00ml} \} \\ &= |M_5|^2 + \text{Re}(M_1 M_2^*), \\ F_2(\vartheta) &= I_{0000} \{ \sin^2(\vartheta/2) A_{00mm} + \cos^2(\vartheta/2) A_{00ll} - \sin(\vartheta) A_{00ml} \} \\ &= \frac{1}{2} \{ |M_3|^2 + |M_4|^2 - |M_1|^2 - |M_2|^2 \}. \end{aligned} \quad (10)$$

Here the observables I_{0000} , A_{00mm} , A_{00nn} , A_{00ll} , A_{00ml} and the helicity amplitudes M_i are defined as in Ref.[6].

Let $\mathbf{P}_T$ be the target polarization vector, and $\boldsymbol{\zeta}_T = \mathbf{P}_T/P_T$. We direct the quantization axis along the unit vector $\boldsymbol{\zeta}_T$. Averaging the cross section in Eq.(9) over particles in the target, we obtain for the quantities $\Omega_{\pm}^{out}$

$$\begin{aligned} \Omega_{\pm}^{out} &= n f \{ \sigma_0^{out} \pm P_T [\sigma_1^{out} + (\boldsymbol{\zeta}_T \cdot \boldsymbol{\nu})^2 (\sigma_2^{out} - \sigma_1^{out})] \}, \\ \sigma_i^{out} &= 2\pi \int_{\theta_{acc}}^{\pi/2} d\vartheta \sin \vartheta F_i(\vartheta). \end{aligned} \quad (11)$$

Here n is an areal density of the target and f is a revolution frequency. The function $P_B(t)$ in Eq.(5) contains $\Omega_{\pm}^{out}$ only in the combination $\Omega_+^{out} - \Omega_-^{out}$. For $|\sigma_2| > |\sigma_1|$, this difference is maximal at $\boldsymbol{\zeta}_T \parallel \boldsymbol{\nu}$. For $|\sigma_2| < |\sigma_1|$, the

difference is maximum at $\zeta_T \perp \nu$. In Eqs.(5) and (6), the quantities Ω_{+-} and Ω_{-+} , which are related to the spin-flip transitions, have the form

$$\begin{aligned}\Omega_{-+} + \Omega_{+-} &= nf \left\{ \sigma_{2s} + (\zeta_T \cdot \nu)^2 (\sigma_{1s} - \sigma_{2s}) \right\}, \\ \Omega_{-+} - \Omega_{+-} &= nf P_T \left\{ \sigma_{2d} + (\zeta_T \cdot \nu)^2 (\sigma_{1d} - \sigma_{2d}) \right\}, \\ \sigma_X &= 2\pi \int_0^{\theta_{acc}} d\vartheta \sin \vartheta G_X(\vartheta).\end{aligned}\tag{12}$$

In terms of the helicity amplitudes, the functions $G_X(\vartheta)$ are

$$\begin{aligned}G_{1s}(\vartheta) &= \frac{1}{2} \left\{ |2 \cos(\vartheta/2) M_5 + \sin(\vartheta/2) (M_1 + M_3)|^2 + |2 \sin(\vartheta/2) M_5 + \cos(\vartheta/2) (M_4 - M_2)|^2 \right. \\ &\quad \left. + \sin^2(\vartheta/2) |M_1 - M_3|^2 + \cos^2(\vartheta/2) |M_2 + M_4|^2 \right\}, \\ G_{2s}(\vartheta) &= \frac{1}{2} \left\{ G_{1s} + \cos^2(\vartheta/2) |M_1 - M_3|^2 + \sin^2(\vartheta/2) |M_2 + M_4|^2 \right\}, \\ G_{1d}(\vartheta) &= \sin \vartheta \operatorname{Re} \left[M_5^* (M_2 + M_4 + M_3 - M_1) \right] + \sin^2(\vartheta/2) (|M_3|^2 - |M_1|^2) \\ &\quad + \cos^2(\vartheta/2) (|M_4|^2 - |M_2|^2), \\ G_{2d}(\vartheta) &= \operatorname{Re} \left\{ \frac{1}{2} \sin \vartheta \left[M_5^* (M_2 + M_4 + M_3 - M_1) \right] + M_2^* (M_1 - M_3) \right. \\ &\quad \left. + \sin^2(\vartheta/2) (M_4^* M_1 + M_2^* M_3) \right\}.\end{aligned}\tag{13}$$

Each amplitude M_i can be represented as a sum $M_i = M_i^{em} + M_i^h$ of a pure electromagnetic amplitude, M_i^{em} , and the hadronic amplitude, M_i^h . The latter also accounts for the electromagnetic interaction. The amplitudes M_i^h are not singular at small scattering angle ϑ . More precisely, at $\vartheta \rightarrow 0$, $M_{1,2,3}^h$ are nonzero constant, $M_4^h \propto \vartheta^2$, and $M_5^h \propto \vartheta$. In the nonrelativistic limit, the amplitudes M_i^{em} pass into the amplitude M_i^C , which are the matrix element of the operator $\hat{M}^C$ in the spin space [7]

$$\begin{aligned}\hat{M}^C &= f(\vartheta) - \frac{1}{2} (1 + \boldsymbol{\sigma}_b \cdot \boldsymbol{\sigma}_t) f(\pi - \vartheta), \\ f(\vartheta) &= -\frac{\alpha}{4vp \sin^2(\vartheta/2)} \exp\{-i(\alpha/v) \ln[\sin(\vartheta/2)]\},\end{aligned}\tag{14}$$

where $v = p/m_p$ is the proton velocity in the center-of-mass frame, $\boldsymbol{\sigma}$ are the Pauli matrices, α is the fine structure constant. The presence of the spin operators in $\hat{M}^C$ is completely due to the identity of protons. From Eq.(14), we obtain for M_i^C

$$\begin{aligned}M_1^C &= \cos^2(\vartheta/2) f(\vartheta) + \sin^2(\vartheta/2) f(\pi - \vartheta), \\ M_2^C &= -[\sin^2(\vartheta/2) f(\vartheta) + \cos^2(\vartheta/2) f(\pi - \vartheta)], \\ M_3^C &= \cos^2(\vartheta/2) [f(\vartheta) - f(\pi - \vartheta)], \\ M_4^C &= \sin^2(\vartheta/2) [f(\vartheta) - f(\pi - \vartheta)], \\ M_5^C &= -\frac{1}{2} \sin \vartheta [f(\vartheta) - f(\pi - \vartheta)].\end{aligned}\tag{15}$$

Using Eq.(15), we can consider the interrelation between the electromagnetic and hadronic contributions to $\Omega_{\pm}^{out}$. If $\theta_{acc} \ll \alpha/(vpH)$, H is a typical magnitude of the hadronic amplitudes, then the main contribution to the cross section σ_0 in Eq.(1) comes from the integration region $\vartheta \sim \theta_{acc} \ll 1$, where $F_0(\vartheta) \simeq |f(\vartheta)|^2$. Thus, this contribution has pure electromagnetic origin:

$$\sigma_0 \approx \sigma_0^C = \pi \alpha^2 / (vp \theta_{acc})^2.\tag{16}$$

For the electromagnetic part of the functions $F_{1,2}$, we have $F_1^C = F_2^C = -\operatorname{Re}[f^*(\vartheta) f(\pi - \vartheta)]$. The corresponding contribution to $\sigma_{1,2}$ reads

$$\sigma_1^C = \sigma_2^C = -\frac{\pi \alpha}{2vp^2} \sin \Psi, \quad \Psi = \frac{\alpha}{v} \ln(2/\theta_{acc}).\tag{17}$$

The interference terms σ_i^{int} in σ_i can be estimated to the logarithmic accuracy as

$$\begin{aligned}\sigma_0^{int} &= -\frac{2\pi}{p} \left\{ \sin \Psi \operatorname{Re}[M_3^h(0) + M_1^h(0)] + (1 - \cos \Psi) \operatorname{Im}[M_3^h(0) + M_1^h(0)] \right\}, \\ \sigma_1^{int} &= -\frac{2\pi}{p} \left\{ \sin \Psi \operatorname{Re} M_2^h(0) + (1 - \cos \Psi) \operatorname{Im} M_2^h(0) \right\}, \\ \sigma_2^{int} &= -\frac{2\pi}{p} \left\{ \sin \Psi \operatorname{Re}[M_3^h(0) - M_1^h(0)] + (1 - \cos \Psi) \operatorname{Im}[M_3^h(0) - M_1^h(0)] \right\}.\end{aligned}\quad (18)$$

We illustrate a scale of different contributions, giving their numerical values corresponding to the parameters of the experiment [3]: $E_{lab} = 23\text{MeV}$, $\theta_{acc} = 8.8\text{mrad}$. Using the data base [8] for the hadronic amplitudes M_i^h , we obtain

$$\begin{aligned}\sigma_0 &= 6444\text{ mb}, & \sigma_0^{int} &= -56\text{ mb}, & \sigma_0^C &= 6357\text{ mb}, \\ \sigma_1 &= -89\text{ mb}, & \sigma_1^{int} &= 39\text{ mb}, & \sigma_1^C &= -1\text{ mb}, \\ \sigma_2 &= -66\text{ mb}, & \sigma_2^{int} &= 66\text{ mb}, & \sigma_2^C &= -1\text{ mb}.\end{aligned}\quad (19)$$

Recollect that $\sigma_i = \sigma_i^{em} + \sigma_i^{int} + \sigma_i^h$ and $\sigma_i^{em} \simeq \sigma_i^C$ in the nonrelativistic case. Note that the numbers obtained from Eqs.(16), (17) and (18) are in good agreement with that in Eq.(19). The role of interference of the hadronic and electromagnetic amplitudes is additionally illustrated by Fig.1. In this figure, the function $2\pi \sin \vartheta F_1(\vartheta)$ calculated with the use of the full amplitudes $M_i = M_i^{em} + M_i^h$ (solid curve) is compared to that obtained using the hadronic amplitudes M_i^h only (dashed curve). Drastic modification of the function $F_1(\vartheta)$ at small ϑ , as compared to the hadronic contribution, is due to interference (the pure electromagnetic contribution is negligible).




FIG. 1: The function $2\pi \sin \vartheta F_1(\vartheta)$ in units mb, calculated with the use of full amplitudes (solid curve) and of hadronic amplitudes only (dashed curve).

Let us consider the quantities σ_{1s} , σ_{2s} , σ_{1d} , and σ_{2d} (see Eq.(12)), which determine the functions Ω_{+-} and Ω_{-+} . The latter correspond to the spin-flip transitions at $\vartheta \leq \theta_{acc} \ll 1$. Using the nonrelativistic electromagnetic amplitudes from Eq.(15), we obtain for the small-angle asymptotics of the functions $G_X(\vartheta)$

$$\begin{aligned}G_{1s} &= |M_2^h(0)|^2, & G_{2s} &= \frac{1}{2}|M_2^h(0)|^2 + \frac{1}{2}|M_1^h(0) - M_3^h(0)|^2, \\ G_{1d} &= -|M_2^h(0)|^2, & G_{2d} &= \operatorname{Re}\{M_2^{h*}(0)[M_1^h(0) - M_3^h(0)]\}.\end{aligned}\quad (20)$$

This contribution is purely hadronic, and the corresponding cross sections can be estimated as $\sigma_s \sim \sigma_d \sim \theta_{acc}^2 \sigma_1^h$, being negligibly small.

If we take into account the first relativistic correction to the electromagnetic amplitudes (see, e.g., [9]), then the additional term appears in the functions G_{1s} and G_{2s} :

$$\delta G_{1s} = 2\delta G_{2s} \simeq \frac{1}{8} \left[\frac{\alpha(4\kappa + 3)}{m_p \vartheta} \right]^2, \quad (21)$$

where $\kappa = 1.79$ is the anomalous magnetic moment of the proton. Though the corresponding cross sections is logarithmically enhanced, it is negligibly small being proportional to α^2/m_p^2 . Thus, scattering events, where projectiles stay in the beam, do not lead to the beam polarization.

IV. PURE ELECTRON TARGET

The time dependence of the beam polarization is described again by Eq.(5), but with another expressions for the cross sections in Eqs.(11) and (12), which correspond now to the pure electromagnetic electron-proton interaction. The cross section accounting for the polarizations of the initial and final proton and electron is well known (see, e.g.,[10]).

The functions Ω_+^{out} and Ω_-^{out} are related to the scattering events, where particles go out from the beam. Below we consider head-on collisions of electrons and protons in the laboratory frame. For any storage ring, $\theta_{acc} \gg m_e/m_p \approx 0.5\text{mrad}$. Then the particle loss occurs at $p_e > p^{out}$, where

$$p^{out} = \frac{p_p \theta_{acc} m_p}{\varepsilon_p + p_p}, \quad (22)$$

$\varepsilon_p = \sqrt{p_p^2 + m_p^2}$ is the energy and p_p is the momentum of a proton, p_e is the momentum of an electron. For $p_e < p_p m_p/(\varepsilon_p + p_p + m_p)$, there is the maximal scattering angle θ_{max} ,

$$\theta_{max} = \arcsin \left[\frac{\varepsilon_p p_e + \varepsilon_e p_p}{m_p(p_p - p_e)} \right], \quad (23)$$

where ε_e is the electron energy. In particular, $\theta_{max} = m_e/m_p$ for $p_e = 0$. For $p_e > p^{out}$, the cross sections σ_0^{out} , σ_1^{out} , and σ_2^{out} in Eq.(11) can be estimated as

$$\sigma_0^{out} \sim \frac{4\pi\alpha^2}{(p_p \theta_{acc})^2} \left(\frac{\varepsilon_e \varepsilon_p + p_e p_p}{\varepsilon_e p_p + \varepsilon_p p_e} \right)^2, \quad \frac{\sigma_1^{out}}{\sigma_0^{out}} \sim \frac{m_e m_p (p_p \theta_{acc})^2}{(\varepsilon_e \varepsilon_p + p_e p_p)^2}, \quad \frac{\sigma_2^{out}}{\sigma_0^{out}} \sim \frac{(p_p \theta_{acc})^2}{\varepsilon_e \varepsilon_p + p_e p_p}. \quad (24)$$

For the time $t \sim \Omega_{tot}^{-1}$, when $N(t)/N(0)$ is not too small (see Eq.(5)), the ratios $\sigma_1^{out}/\sigma_0^{out}$ and $\sigma_2^{out}/\sigma_0^{out}$ give the estimations of the beam polarization $P_B(t)$ for $\zeta_T \perp \nu$ and $\zeta_T \parallel \nu$, respectively. These ratios are maximal for $p_e \gtrsim p^{out}$, so that $P_B < m_e/m_p$ for $\zeta_T \perp \nu$, and $P_B < p_p \theta_{acc}/m_p$ for $\zeta_T \parallel \nu$. In both cases P_B is too small, and the mechanism using the loss of particles due to proton scattering on polarized electrons (filtering) does not work for any parameters of the electron beam.

Let us consider now the mechanism of the polarization buildup without loss of the particles, which is due to the spin-flip transitions. Starting with the general expressions for the cross section of polarized electron-proton scattering, [10], we find that the cross sections σ_s and σ_d in Eq.(12) are maximal at small relative velocity of electron and proton. In this case, we estimate $\sigma_d \sim 16\pi\mu_p^2 \sim 10^{-3}\text{mb}$, μ_p is the proton magnetic moment. As compared with σ_d , the cross section σ_s is enhanced by some logarithmic factor, which can not change essentially the conclusion on its smallness. We emphasize that the cross sections are small because the spin-flip amplitude does not contain a term $\propto 1/\vartheta^2$ at $\vartheta \rightarrow 0$, contrary to the amplitude without the spin-flip transition where such term (Coulomb term) is present. Thus, for a pure electron target, the mechanism of polarization buildup based on the spin-flip transitions also does not work.

In conclusion, it is demonstrated that, for $\theta < \theta_{acc}$ (a proton remains in the beam), the polarization buildup is completely due to the spin-flip transitions. The corresponding cross sections turn out to be negligibly small for both proton-proton and proton-electron scattering. For a pure electron target, filtering mechanism also does not provide a noticeable polarization. Evidently, these statements are valid for the antiproton beam as well. Thus, the filtering method using a hydrogen gas target with the proton polarization seems to be the most promising way to polarize stored antiprotons.

ACKNOWLEDGMENTS

We are grateful to I.A.Koop, V.V.Parkhomchuk, Yu.M.Shatunov, A.N.Skrinsky, and D.K.Toporkov for stimulating discussions. This work was supported in part by RFBR Grants 03-02-16510 and 05-02-16079.

[1] F. Rathmann et al, Phys. Rev. Lett. **94**, 014801 (2005).

- [2] P.L.Csonka, NIM **63**, 247 (1968).
- [3] F. Rathmann et al, Phys. Rev. Lett. **71**, 1379 (1993).
- [4] H.O. Meyer, Phys. Rev. E **50**, 1485 (1994).
- [5] C.J. Horowitz and H.O. Meyer, Phys. Rev. Lett. **72**, 3981 (1994).
- [6] J. Bystricky, F. Lehar, and P. Winternitz, J. Phys. (Paris) **39**, 1 (1978).
- [7] L.D. Landau, E.M. Lifshits, *Quantum Mechanics, Nonrelativistic Theory*, Pergamon, Oxford (1965).
- [8] Nijmegen data base, accessible via <http://nn-online.org>
- [9] V.B. Berestetski, E.M. Lifshits, and L.P. Pitaevsky, *Quantum Electrodynamics*, Pergamon, Oxford (1982).
- [10] J.H. Scofield, Phys. Rev. **141**, 1352 (1966).